

Mallet Injury

Hand Service

What is a mallet finger (thumb) injury?

A mallet finger injury is usually caused when the end joint of a finger is forced to bend forwards suddenly. Causing the tendon that straightens the tip of the finger to tear or pull away from bone.

Symptoms:

- Inability to straighten the end joint of the finger.
- Pain & swelling may be present.

Treatment:

A splint will be applied to your finger to keep the end joint straight while the tendon heals. Initially, you will be provided with a temporary 'off the shelf' splint which will adequately hold your finger straight. However, most patients with this injury will be referred to the Hand Therapy Department for a more bespoke, comfortable splint.

Constant Splinting Phase:


- The splint is worn **constantly** day & night for a minimum of **six to eight** weeks; your Consultant or Therapist will advise how long you will need to wear the splint.
- It is very important that the splint is always securely taped or strapped to your finger & that the end joint **DOES NOT BEND** during this period of splinting. As any movement of the end finger joint will delay healing & may cause a poor outcome.

Take all precautions to keep the splint & finger dry at all times to prevent your skin from becoming very sore.

It is advisable to remove the splint at least once a week to clean your finger & splint, it is much easier if someone assists you with this. Some people may want to do this daily, this is perfectly acceptable. To clean your finger, place the tip of your finger (palm side) lightly on to the tabletop. Remove the tape / straps and carefully remove the splint. Using a Wet Wipe carefully clean your finger while it is resting on the tabletop, dry, then reapply the splint securely. Please ensuring while cleaning your finger you do not allow the end joint to bend as this could cause further damage.

Your Hand Therapist will demonstrate how to fit & remove the splint & clean your finger.

Pictured is the most frequently supplied splint.



Protective splinting phase:

After six or eight weeks your Hand Therapist will check your finger and if the tendon is sufficiently healed you will progress onto a two to four week period of protective splinting. You are still required to wear the splint overnight & during the day, outdoors, at school, at work & during moderate activities. However, it can be removed in a safe environment, also for cleaning your hand & to perform gentle exercises as demonstrated by your Hand Therapist. It takes twelve weeks for the tendon to fully heal. Your finger will feel stiff and movement will improve with the exercises, however please do not be tempted to physically push your finger into a bent position.

Contact your Therapist if your splint causes any of the following problems:

- Increased pain
- Skin irritation/soreness; you think the splint is rubbing
- You think your finger has become more swollen
- If your splint feels loose when swelling reduces

If your hand or fingers turn blue, feel cold, or tingle:

- Check the straps are not too tight
- Loosen the straps and seek medical advice if the symptoms continue.

If problems arise, please contact the Hand Therapy Department to arrange an appointment. We can only see you if you have been referred by your GP, Consultant or Walk in Centre / A&E. In an emergency, seek medical advice from one of the above.

Name of Therapist..... Date:.....

Other Information:

Contact Details: Ashford & St Peters Hand Therapy Department

Tel: 01784 88 4043

Opening Hours: Monday to Friday 8am to 4pm

Reference: British Society for Surgery of the Hand. 2016 Mallet Finger Injury, leaflet, www.bssh.ac.uk

Version: 2

Published: January 2023

Review: January 2025

We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.



اگر نیاز بہ ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید۔
 ने सुवातुं उरनामे दी लेऊ वै उं विरथा बरवे इस नंबर से हें बरवे: 01932 723553
 اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براؤزر میں اس فون نمبر 01932 723553 پر رابطہ کریں
 Se precisa de uma tradução por favor contacte: 01932 723553

आपनार अनुवादर दरकार हले एखाने योगायोग करुन : 01932 723553
 यदि आपको अनुवाद की जरूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553
 Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

Ashford Hospital London Road, Ashford, Middlesex, TW15 3AA Tel: **01784 884488**

St. Peter's Hospital Guildford Road, Chertsey, Surrey, KT16 0PZ Tel: **01932 872000**

Website: www.ashfordstpeters.nhs.uk