



We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.

To use the Text Relay service, prefix all numbers with 18001.

اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید۔

ਜੇ ਤੁਹਾਨੂੰ ਤਰਜਮੇ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ ਤੇ ਫੋਨ ਕਰੋ: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براہ کرم اس فون نمبر 01932 723553 پر رابطہ کریں

Se precisa de uma tradução por favor contacte: 01932 723553

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553

यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

Ashford Hospital
London Road
Ashford, Middlesex
TW15 3AA
Tel: **01784 884488**

St. Peter's Hospital
Guildford Road
Chertsey, Surrey
KT16 0PZ.
Tel: **01932 872000**

Website: www.ashfordstpeters.nhs.uk

Information for Women Taking Aspirin in Pregnancy Maternity Department

Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty.

If they cannot resolve your concern, please contact our Patient Experience Team on 01932 723553 or email asp-tr.patient.advice@nhs.net. If you remain concerned, the team can also advise upon how to make a formal complaint.

Author: Dr Sian McDonnell

Department: Obstetrics

Version: 4

Published: December 2020

Review: June 2022

Can I still take Aspirin if I am taking other medication?

Aspirin can occasionally interact with other medications so it is important that you tell the midwife or doctor if you are taking any other medications or herbal remedies, including those bought over the counter without a prescription.

For more information, speak with your midwife or GP or phone **01932 722389 or 722388**

You have been given this leaflet because you have been advised to take Aspirin during your pregnancy.

When the placenta, which feeds your baby and helps your baby, does not work properly it can lead to your baby being smaller and needing early delivery. It could also lead to you developing pre-eclampsia. Very rarely, the placenta can cause the baby to be stillborn or die within the first few months of life. Taking Aspirin has been shown to help reduce the risk of these problems.

Why am I at higher risk?

At your first few antenatal appointments with the midwife or doctor, they will have asked you questions to help identify if you are at higher risk of problems with the placenta. These include:

- This being your first pregnancy
- You being 40 or over
- Your last pregnancy being more than 10 years ago
- Your BMI (body mass index) being over 35
- You are expecting more than one baby this pregnancy
- Blood pressure before you became pregnant
- Raised blood pressure or pre-eclampsia in a previous pregnancy
- A previous small baby, either <2.5kg or <10th centile
- Chronic kidney disease or diabetes.
- Low Pregnancy Associated Plasma Protein A – Papp-A <0.41MoM

Signs and symptoms of pre-eclampsia

Pre-eclampsia can cause lots of different symptoms including:

- Headaches
- Vision (eye) problems
- Tummy pain
- Swollen feet, hands and face

Often, there are no symptoms and women feel completely well. This is the reason the midwife or doctor will check your blood pressure and urine each time you see them. It affects as many as 1 in 10 pregnancies but is often mild. In severe cases it can affect 1 in 100 pregnancies.

How will my baby's growth be checked?

Additional scans will be arranged after your 20 week anomaly scan.

Exactly when and how many scans you have depends on your risks and blood flow tests performed at the scan. Your baby does not need to be delivered early just because you are taking Aspirin.

How do I take Aspirin?

Your midwife will tell you at your booking appointment if you should take Aspirin. You will:

- Receive a letter detailing why you have been advised to take Aspirin in your pregnancy
- Take this letter to your first trimester ultrasound scan and the midwives in Antenatal Clinic will give you a 28 day supply
- The same letter then needs to be given to your GP to arrange ongoing supply

It has been shown to be most effective to take 150mg Aspirin (2 'baby' Aspirin) every night from 12 to 36 weeks of pregnancy. Taking it at night has been shown to help the medicine be absorbed.

If you miss the occasional dose, do not worry. Take the dose when you remember but do not take an 'extra' dose.

What are the side effects of Aspirin?

Aspirin is very safe for most women to take.

If you have previously had stomach ulcers, bleeding disorders or asthma, or are allergic to Aspirin, please consult your Obstetrician before taking Aspirin.

It has not been shown to increase the risk of having a miscarriage or bleeding during pregnancy or at the time of delivery. It has not been shown to cause any abnormalities in babies.