



We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.

To use the Text Relay service, prefix all numbers with 18001.

اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید۔

ਜੇ ਤੁਹਾਨੂੰ ਤਰਜਮੇ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ ਤੇ ਫੋਨ ਕਰੋ: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براہ کرم اس فون نمبر 01932 723553 پر رابطہ کریں

Se precisa de uma tradução por favor contacte: 01932 723553

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553

यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

**Ashford Hospital**  
London Road  
Ashford, Middlesex  
TW15 3AA  
Tel: **01784 884488**

**St. Peter's Hospital**  
Guildford Road  
Chertsey, Surrey  
KT16 0PZ.  
Tel: **01932 872000**

Website: [www.ashfordstpeters.nhs.uk](http://www.ashfordstpeters.nhs.uk)

# Mallet Finger

## Paediatric A&E Department




## Introduction and explanation of condition

Your child has torn the tendon that straightens the end joint of their finger. We call this a "mallet injury".

To treat this, the joint must be held straight in a splint for 6 - 8 weeks as tendons heal very slowly. Your child will be seen in the fracture clinic during this time.

If your child's finger is allowed to bend during this time, healing will not occur and the "droop" of the finger will be permanent. The splint should, therefore, be removed **ONLY** for cleaning and **while keeping the finger straight**.

## The treatment



The fingertip droops because the tendon to extend it has snapped.

The treatment is to avoid letting the fingertip drop **at all times** for a minimum of 6 – 8 weeks. Your child must wear the splint at all times except for cleaning (see above)



The splint (which may vary in design) will allow the tip to be kept extended when the hand is in use.

## Do Not

- Attempt to examine the healing by bending the fingertip until advised by the treating doctor.

## Do

- Ensure your child's hand is elevated in a sling for the first week and that they move all the joints normally except the splinted one.
- Make sure that you have follow up appointments at the Fracture Clinic.

## Further Information

Paediatric Accident and Emergency Department  
St. Peter's Hospital, Guildford Road  
Chertsey  
Surrey KT16 0PZ

Telephone: **01932 872000**

For reassurance and advice please contact NHS Direct on: **111**

The following website gives useful advice: [www.patient.co.uk](http://www.patient.co.uk)

---

## Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty.

If they cannot resolve your concern, please contact our Patient Experience Team on 01932 723553 or email [asp-tr.patient.advice@nhs.net](mailto:asp-tr.patient.advice@nhs.net). If you remain concerned, the team can also advise upon how to make a formal complaint.

---

Author: Dr Erin Dawson

Department: Paediatric A&E

Version: 6

Published: Apr 2018

Review: Apr 2020