

Basal Rate Testing

Children and Young Person's Diabetes Service

What is it and why do we do it?

Documenting blood sugar levels 2 hourly for a certain time period in a day without influences from food and bolus insulin and other factors, ascertains if your basal rates are set correctly. As children's bodies grow and develop, basal requirements change and therefore, basal rate testing should be completed several times a year.

Before you start, you need to check:

- BSL in target range 4-10mmol
- Not had recent correction
- Not had hypo in last eight hours
- Last food insulin 4 hours previous therefore no active insulin
- Last food was not high fat or low GI
- No strenuous activity in last 24 hours
- You are not sick or have other external factors that might influence
- You are not due a set change
- Your cannula site is free from lumps
- You are prepared to fast/ have carb free food and drinks for the time period

Basal rate testing times:

- Divide the day into different periods - Overnight, Morning, Afternoon, Evening.
- Only do one period at a time. Generally, start with overnight first.
- At the start of the test for each time period, check BSL and document or add notes on Libre.
- Continue to check 2 hourly until the time period has ended.
- Stop the test if you need a correction or have a hypo
- All food and drink consumed during the test period should be carb free
- **Re-do the time period test again before making any changes.**

Interpreting the results:

- If basal rates are correct then there should be no greater change than 2mmol/L
- If there is a rise of more than 2mmol/L then identify the time of the higher blood sugar and increase the basal by 1 to 2 increments two hours prior to this.
- If there is a decrease of more than 2mmol/L then identify the time the blood sugar was lower and decrease the basal by 1 to 2 increments two hours prior to this.

Test 1:

time	12pm	1am	2am	3am	4am	5am	6am	7am	8am	9am	10am	11am	12am	1pm	2pm	3pm	4pm	5pm	6pm	7pm	8pm	9pm	10pm	11pm	
current basal rate																									
mark when last had insulin and food																									
document BSL at start/end and every 2 hours of test																									

Test 2:

time	12pm	1am	2am	3am	4am	5am	6am	7am	8am	9am	10am	11am	12am	1pm	2pm	3pm	4pm	5pm	6pm	7pm	8pm	9pm	10pm	11pm	
current basal rate																									
mark when last had insulin and food																									
document BSL at start/end and every 2 hours of test																									

Examples of carbohydrate-free foods:

Protein foods such as:

- eggs,
- cheese
- meat, chicken, fish, (no breadcrumb or batter coating)

Low carb vegetables such as:

- tomatoes
- green leafy vegetables
- cucumber
- courgettes
- sugar free drinks
- sugar-free jelly

Contact: asp-tr.generalpaeddiabetes@nhs.net

Reference: Medtronic resources

Author: Claire Berriman

Dept: Paediatric Diabetes

Version: 1

Last Review: Nov 2022

Next Review: Nov 2024

We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.



اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید۔

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553

ਜੇ ਤੁਹਾਨੂੰ ਤਰਜਮੇ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ ਤੇ ਫੋਨ ਕਰੋ: 01932 723553

यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براؤزر اس فون نمبر 01932 723553 پر رابطہ کریں

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

Se precisa de uma tradução por favor contacte: 01932 723553

Ashford Hospital London Road, Ashford, Middlesex, TW15 3AA Tel: 01784 884488

St. Peter's Hospital Guildford Road, Chertsey, Surrey, KT16 0PZ Tel: 01932 872000

Website: www.ashfordstpeters.nhs.uk