



We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.

To use the Text Relay service, prefix all numbers with 18001.

اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید۔

ਜੇ ਤੁਹਾਨੂੰ ਤਰਜਮੇ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ ਤੇ فون ਕਰੋ: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براؤکرم اس فون نمبر 01932 723553 پر رابطہ کریں

Se precisa de uma tradução por favor contacte: 01932 723553

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553

यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

Ashford Hospital
London Road
Ashford, Middlesex
TW15 3AA
Tel: **01784 884488**

St. Peter's Hospital
Guildford Road
Chertsey, Surrey
KT16 0PZ.
Tel: **01932 872000**

Website: www.ashfordstpeters.nhs.uk

Pain Management Service: After your treatment



Pain Management Services

Introduction

You have attended Ashford and St Peter's Hospitals NHS Foundation Trust to have an injection treatment to help your pain. This leaflet aims to give you support and advice following your procedure.

What can I expect to feel after my procedure? Immediately after the procedure

Chronic pain is complex and not everyone will experience exactly the same sensations or feelings after a procedure.

Immediately after your procedure you may have numbness around the treatment area. This may sometimes be accompanied by weakness or heaviness. This is due to the use of local anaesthetic and this will wear off in a variable length of time, usually a few hours.

If you have significant sensitivity you may feel pain at the injection sites in spite of the local anaesthetic. This may sometimes be associated with stiffness or muscle tension. If you are feeling this please tell the nurse looking after you on day ward and you may take some of your normal pain medicines, or if required an additional pain medicine can be prescribed for you.

Occasionally you may feel sick or dizzy. Please tell the nurse on day ward if this is the case and you will have a check of your

Additional useful information in respect of the Chronic Pain Service at Ashford and St Peter's may be obtained by logging on to the hospital website www.ashfordstpeters.nhs.uk.

From the drop down menu select:

- A-Z of Departments - Chronic Pain Management

Leaflets

For patient information leaflets - Pain management (Chronic)
<http://www.asph.nhs.uk/leaflets-pain>
<https://www.ashfordstpeters.info/leaflets>

Alternatively the British Pain Society provides a comprehensive website which you may find useful www.britishpainsociety.org

Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty.

If they cannot resolve your concern, please contact our Patient Experience Team on 01932 723553 or email asp-tr.patient.advice@nhs.net. If you remain concerned, the team can also advise upon how to make a formal complaint.

Author: Dr Rose Block

Version: 2

Department: Pain Management Services

Published: August 2021

Review: August 2023

Please make gradual changes only and do not stop your medication suddenly. It is best to make only one medication change at a time when managing chronic pain, and to then allow a period of stability before initiating further changes.

An appointment for follow up may be arranged in pain clinic.

If you have been given a graph to chart your pain level after treatment please bring this to your review appointment, or post it to the department prior to review.

Further Information

If you have any questions regarding treatment in the Chronic Pain Service, please call our Pain Clinic Advice Line on **01784 884970**.

For questions regarding your appointment or any other administrative matter, please call the Surrey iMSK line: **01932 722730, select option 5 for procedure booking office or option 4 for pain clinic office.**

Website: www.surreyimsk.com

observations. This may include blood pressure and pulse rate. It may include blood sugar measurement. In most cases this will resolve rapidly without specific treatment. Occasionally you may need to have fluids prescribed, or medication, or be further reviewed by your doctor. Very rarely you may need to stay a little longer on day ward for observation.

Please ensure you are driven home by a responsible adult who will be able to ensure you are settled at home after your procedure.

At home after my procedure

After the local anaesthetic wears off you may experience an increase of pain. In almost all cases this will respond to regular use of your normal pain medicines. You should take these regularly as prescribed.

We would advise that you do not apply heat to the area on the first day as this will encourage bruising. Application of cold packs may be helpful.

Activity

In general it is important to maintain mobility. If you do have pain immediately following the procedure it may be advisable to reduce activity. However gentle paced mobility and muscle stretches should be maintained throughout.

Dressing

Please keep any dressings on for 24 hours after which you may soak it off in a shower or bath.

If you have had sacroiliac joint radiofrequency treatment you may have steristrips and a dressing – please keep this on for 5-7 days before removing these at home.

You may notice some bruising at the injection site. This will resolve in a few days.

Steroids are frequently used as part of the injection therapy. If you have diabetes then the steroid may sometimes alter blood sugar levels. If this occurs please contact your diabetic team at your local surgery or at the Trust for advice.

Steroids may sometimes cause irregularity of menstrual cycle, for up to 3 months.

If you are of childbearing age, we advise you to avoid conception of pregnancy for 3 months following treatment, and to use effective contraception.

Rarely, you may feel unwell after a procedure. If this is associated with fever and / or swelling and redness at the injection site please seek medical advice to exclude the rare possibility of infection.

Rarely, you may develop more unusual side effects. If this occurs please contact your GP or contact the pain clinic advice line for non-urgent advice.

Rarely, people feel unwell and do not feel able to await an appointment with the GP or a call back from the advice line, if this occurs please attend A and E.

Later after my procedure

Many procedures use depot steroid and this will take 2 weeks to be fully effective.

Most of the benefit you feel will be expected to be in the treated area, benefit for other pains are possible but less likely.

Some patients may experience flare up of their pain after injection treatment, in the majority of cases this will settle in a few days with normal pain medicines, and pacing of activity.

Some patients may experience flare up of other pain related to the position required to do the treatment procedure, this will also resolve with simple measures.

Radiofrequency denervation for spinal pain may cause a longer period of flare up, in the majority of cases this will be managed with normal pain medicines and pacing of activity.

You will gain most benefit from your procedure if you are able to enhance your activity levels.

Please maintain your normal pain medication regime after your procedure. When you have had time to fully recover from your procedure you may wish to consider changes to your medication. In the first instance please discuss this with your GP.