



We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.

You can use Text Relay to turn telephone communications into text.
Use the Relay UK app on your phone, tablet or PC.
You can also use Relay UK via Minicom or Uniphone.

اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید.

ने सुगठु उत्तमे ची लेउ वै उं विरथा वरवे इस निसर उे देन वरवे: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براہ کرم اس فون نمبر 01932 723553 پر رابطہ کریں

Se precisa de uma tradução por favor contacte: 01932 723553

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553

यदि आपको अनुवाद की जरूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

Ashford Hospital
London Road
Ashford, Middlesex
TW15 3AA
Tel: **01784 884488**

St. Peter's Hospital
Guildford Road
Chertsey, Surrey
KT16 0PZ.
Tel: **01932 872000**

Website: www.ashfordstpeters.nhs.uk

Exercise and Insulin

Nutrition and Dietetics

Introduction

Effects of Exercise on Blood Glucose Levels

Different types of exercise may affect your blood glucose levels in different ways. Testing blood glucose levels before, during and after exercise will help you identify how **your** body responds and how best to manage the effects.

Low blood glucose levels may be caused by:

- Increased uptake of glucose into the exercising muscles to provide energy during and after exercise
- Faster insulin absorption due to increased blood flow-avoid injecting into exercising muscles
- Too much circulating insulin / insufficient carbohydrate ingestion
- Increased insulin sensitivity for up to 24 hours post exercise - **increased risk of hypoglycaemia for all types of exercise.**

Increased blood glucose levels may be caused by:

- Too little circulating insulin to allow glucose uptake into muscle cells
- Hormone responses increase glucose production from the liver

Useful websites

Runsweet

<http://www.runsweet.com/>

Diabetes UK

<https://www.diabetes.org.uk/>

American Diabetes Association

<https://diabetes.org/>

Dietitian:

Contact details:

Patient name:

Date:.....

Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty.

If they cannot resolve your concern, please contact our Patient Experience Team on 01932 723553 or email asp-tr.patient.advice@nhs.net. If you remain concerned, the team can also advise upon how to make a formal complaint.

Author: Diabetes Dietitian

Department: Nutrition & Dietetics

Version: 5

Published: Dec 2023

Review: Dec 2025

Unplanned exercise - carbohydrate supplementation and diet

Unplanned exercise may need additional carbohydrate intake.
Consider:

- Pre exercise blood glucose level
- When was your last meal / insulin bolus?
- Type / duration of exercise

Post exercise

There is an increased risk of hypoglycaemia for up to 24 hours post exercise. Monitoring blood glucose levels and noting trends in glucose fluctuations will help you avoid this. You could:

- Reduce your post exercise meal bolus
- Have a carbohydrate rich snack after exercise (without extra insulin), particularly if exercising in the evening

General safety

- Always ensure you have some fast-acting carbohydrate with you in the event of hypoglycaemia
- Remember to take your blood glucose monitor with you
- Carry some form of diabetes identification and, if possible, let other people know how long you will be
- Wear appropriate footwear
- Stay hydrated

Type and Intensity of Exercise

Aerobic

Longer duration, moderate intensity endurance (more than 30 minutes) e.g. brisk walking, running, swimming, cycling.

- Greater risk of hypoglycaemia during and after exercise

Anaerobic

Short duration, high intensity (no more than a few minutes) e.g. sprinting, weightlifting.

- Greater risk of hyperglycaemia during and immediately after exercise

Intermittent high intensity

Lots of sports have periods of low or moderate intensity activity punctuated by bursts of high intensity e.g. football, netball.

- Less immediate risk of hypoglycaemia, possible hyperglycaemia

Variables which can cause hypoglycaemia

- Repeated episodes of exercise
- Recent hypoglycaemia (within last 24 hours)
- **NB. Do not exercise within 24 hours of an episode of severe hypoglycaemia**
- Changes in weather temperature
- Altitude

Variables which can cause hyperglycaemia

- Competition stress
- Heat stress including humidity
- Dehydration
- High blood glucose pre-exercise

Blood glucose (BG) levels at the start of exercise

BG mmol/l	Action
<7 mmol	Consider having some carbohydrate
7-12 mmol	Exercise may be started
>12-16 mmol (no ketosis)	Delay carbohydrate during exercise until BG reduces. Mild to moderate aerobic exercise e.g. light jogging / walking may commence with regular BG monitoring
>16 mmol (no ketosis)	Delay exercise until BG falls to safe level especially if planning intense exercise - BG may rise further
Ketosis	Delay exercise until ketones cleared

BG monitoring

- Before exercise
- Every 20-30 minutes during
- At the end of exercise
- Ideally, later on as well

Use this information to decide whether changes in insulin doses or carbohydrate intake are necessary.

Carbohydrate intake

*Before exercise

Ensure you have had some moderate GI (Glycaemic Index) carbohydrate 1-2 hours before e.g. hummus and crispbreads, banana, and almonds.

During

For moderate intensity endurance exercise, quickly digested (high GI) carbohydrate may be required after 20 minutes of exercise in an amount of 30-60g spread across an hour. Have fluids- dehydration leads to poor performance.

Lower intensity or intermittent high intensity exercise may need smaller quantities or none at all.

Immediately after exercise

Foods / drinks containing high GI carbohydrates should be available as soon as possible after exercise. Proper rehydration is essential.

Your dietitian will be able to help you with this and provide information about the Glycaemic Index.

*Planned exercise - insulin dose adjustment

If you are undertaking moderate intensity endurance exercise within 90 minutes of a meal, consider reducing the meal insulin bolus. As people respond differently to exercise, it is not possible to give exact guidelines, but it is best to start with a small reduction, e.g. 20%, and monitor results. Discuss changes to your basal insulin with your healthcare professional.