

# Thumb Carpometacarpal Joint (CMCJ) Replacement Hand Therapy Department

## The Surgery

The trapezium bone sits at the bottom of the thumb to form the 'first CMC joint'. This bone is very prone to wear and tear which can cause pain and limit hand function. There are different surgeries to address this problem each with a different post-operative rehabilitation plan.

The surgery that you have had involves removing the first arthritic joint at the bottom of your thumb and replacing it with an artificial joint also called an implant / prosthesis. Thumb CMCJ prosthetic replacement aims to restore thumb biomechanics and improve pain and function. There are various types of implants that can be used based on your condition and surgeon's preference.



MAIA implant



Touch implant

## Post-Operative care

Immediately Following Surgery

- You will be in a bulky dressing made more rigid with plaster
- This will remain in situ for 2 weeks

## Two Weeks Following Surgery

You will be seen in Hand Therapy for:

- Fabrication of a thermoplastic splint which will support your thumb and wrist in a comfortable and appropriate position. You will be advised on constant use of the splint, removing only for exercises.
- The range of movement of your joints particularly your thumb and wrist will be assessed, and you will be advised a home exercise programme comprising of (but, not exclusive to) the following gentle active exercises:
  - Wrist flexion / extension
  - Thumb IP joint flexion / extension
  - Thumb opposition
  - Thumb circumduction
- You will be advised on scar management once the wound is fully healed.



### **Three Weeks Following Surgery**

You will attend hand therapy for:

- Scar management as appropriate
- Review home exercise programme
- Review splint - continue constant use
- Fine motor activity

### **Four Weeks Following Surgery**

You will attend hand therapy for:

- Continued scar management and advice
- Review splint - continue constant use
- You may commence gentle wrist strengthening as advised by your therapist

Although, at this point in time you may feel that the hand has recovered well from surgery, it remains important to proceed with caution and as advised by your therapist so that you don't put the prosthesis at risk.

### **Six Weeks Following Surgery**


You will attend hand therapy for:

- Advice on removing the splint during the day- you will need to continue with night and protective use
- Advice on using the hand for light activities, e.g. toothbrush, cutlery, light cup of tea
- Assessment of gross grip strength
- Assessment of pinch strength
- Progression of home exercises to include graded hand and thumb strengthening
- Continued scar management and advice

### **Six to Twelve Weeks Following Surgery**

Hand therapy input will continue if there is a specific clinical need and hand therapy programme progressed accordingly. You are likely to have a review with your surgeon to check on your progress.

**Please note - these are just guidelines, and all patients are treated individually depending on their presentation on a case-by-case basis and as guided by the Consultant.**



## Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty. If they cannot resolve your concern, please contact our Patient Experience Team on 01932 723553 or email [asp-tr.patient.advice@nhs.net](mailto:asp-tr.patient.advice@nhs.net). If you remain concerned, the team can also advise upon how to make a formal complaint.

**Author:** Neha Endlay, Specialist Occupational Therapist

**Department:** Hand Therapy

**Version:** 1

**Published:** February 2024

**Review:** February 2026

**We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.**



You can use Text Relay to turn telephone communications into text. Use the Relay UK app on your phone, tablet or PC. You can also use Relay UK via Minicom or Uniphone.

اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید.

आपनार अनुवादर दरकार हले एखाने योगायोग करुन : 01932 723553

ਜੇ ਤੁਹਾਨੂੰ ਤਰਜਮੇ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ ਤੇ ਫੋਨ ਕਰੋ: 01932 723553

यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براؤزر میں اس فون نمبر 01932 723553 پر رابطہ کریں

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

Se precisa de uma tradução por favor contacte: 01932 723553

**Ashford Hospital** London Road, Ashford, Middlesex, TW15 3AA Tel: **01784 884488**

**St. Peter's Hospital** Guildford Road, Chertsey, Surrey, KT16 0PZ Tel: **01932 872000**

**Website:** [www.ashfordstpeters.nhs.uk](http://www.ashfordstpeters.nhs.uk)