



We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.

You can use Text Relay to turn telephone communications into text.  
Use the Relay UK app on your phone, tablet or PC.  
You can also use Relay UK via Minicom or Uniphone.

اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید.

ਜੇ ਤੁਹਾਨੂੰ ਸਹਾਇਤਾ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ 'ਤੇ ਫੋਨ ਕਰੋ: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براہ کرم اس فون نمبر 01932 723553 پر رابطہ کریں

Se precisa de uma tradução por favor contacte: 01932 723553

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553

यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

**Ashford Hospital**  
London Road  
Ashford, Middlesex  
TW15 3AA  
Tel: **01784 884488**

**St. Peter's Hospital**  
Guildford Road  
Chertsey, Surrey  
KT16 0PZ.  
Tel: **01932 872000**

Website: [www.ashfordstpeters.nhs.uk](http://www.ashfordstpeters.nhs.uk)

# Inferior Vena Cava Filter Placement Imaging Department

couple of hours, and you will be able to go home. Take it easy for the rest of the day but you can resume normal activities the next day.

## How long will the filter stay in?

Modern IVC filters can be left in permanently; however, it is becoming more common for these devices to be a temporary solution and removed when they are no longer required. This is often at three months but may occasionally be longer.

## Finally

Some of your questions should have been answered by this leaflet but remember that this is only a starting point for discussion about your treatment with the doctors looking after you. Make sure you are satisfied that you have received enough information about the procedure.

---

## Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty.

If they cannot resolve your concern, please contact our Patient Experience Team on 01932 723553 or email [asp-tr.patient.advice@nhs.net](mailto:asp-tr.patient.advice@nhs.net). If you remain concerned, the team can also advise upon how to make a formal complaint.

---

Author: Mark Mallari

Department: Imaging

Version: 3

Published: Jan 2024

Review: Jan 2027

will be asked to get undressed and put on a hospital gown. You will lie on the X-ray table, generally flat on your back. A small cannula (thin tube) will be placed into a vein in your arm. This is so that a sedative or painkillers can be given if required. You may have monitoring devices attached to your chest and finger and may be given oxygen. The procedure is performed under sterile conditions and the interventional radiologist and radiology nurse will wear sterile gowns to carry out the procedure. The skin near the point of insertion, usually the neck but occasionally the groin, will be swabbed with antiseptic and you will be covered with sterile drapes. The skin and deeper tissues over the vein will be numbed with local anaesthetic. A fine tube (catheter) will be inserted and guided, using the X-ray equipment, into the correct position. Small amounts of dye (contrast agent) are used to check the position of the catheter. The filter is passed through the tube to the exact site and released. Small hooks grip the wall of the vein and stop it moving away.

### **How long will it take?**

Every patient is different, and it is not always easy to predict; however, expect to be in the radiology department for about an hour.

### **What happens afterwards?**

You will be taken back to your ward. Nursing staff will carry out routine observations including pulse and blood pressure and will also check the treatment site. You will generally stay in bed for a

## **Inferior vena cava filter placement**

### **Patient information**

*Reproduced with the kind permission of the British Society of Interventional Radiology (BSIR) 2011*

This leaflet tells you about having an inferior vena cava (IVC) filter inserted. It explains what is involved and what the possible risks are. It is not meant to replace informed discussion between you and your doctor but can act as a starting point for such discussions. If you have any questions about the procedure, please ask the doctor who has referred you or the department which is going to perform it.

### **What is an IVC filter?**

An IVC filter is a small metal device usually placed in a large vein called the inferior vena cava (IVC) that drains blood from the legs and lower part of the abdomen. The IVC filter allows blood to flow through normally but traps any large blood clots, stopping them from getting to your lungs.

### **What is it used for?**

Blood clots (thrombosis) sometimes form in the veins of the legs and pelvis. They are known as a deep vein thrombosis (DVT). The clot can sometimes break free and enter with the blood flow in the lungs and make you very sick. This is called pulmonary embolism

(PE). They can be fatal. An IVC filter prevents a large PE by trapping a clot before it reaches the lungs.

### **Who should have an IVC filter?**

The usual treatment for DVT and PE is drug treatment to thin the blood. This is usually with warfarin. In a few patients, warfarin does not prevent further PEs, in others thinning the blood is too risky. When this happens, patients are considered for treatment by inserting an IVC filter. Very occasionally, a patient is advised to have an IVC filter inserted even though they do not have a DVT or PE at that time. Your doctors will explain the reasons why they think you should have an IVC filter.

### **Are there any risks?**

IVC filter insertion is a very safe procedure. Serious complications are very rare. There may be a small bruise at the needle site and very rarely there may be damage to the vein or blockage of the inferior vena cava. Extremely rarely, the filter can migrate which may require a further procedure to reposition the IVC filter. If you need a magnetic resonance (MRI) scan in the future, you should tell the person doing the scan that you have a filter.

### **Are you required to make any special preparations?**

Insertion of an IVC filter is usually carried out as a day case procedure under local anaesthetic. You may be asked not to eat for six hours before the procedure, although you may still drink clear fluids such as water. If you have any allergies or have previously had a reaction to the dye (contrast agent). You *must* tell the radiology staff before you have the test.

### **Who will you see?**

A specially trained team led by an interventional radiologist within the radiology department. Interventional radiologists have special expertise in reading the images and using imaging to guide catheters and wires to aid diagnosis and treatment.

### **Where will the procedure take place?**

In the interventional suite or theatre, this is usually located within the radiology department. This is similar to an operating theatre into which specialized X-ray equipment has been installed.

### **What happens during the insertion?**

The interventional radiologist will explain the procedure and ask you to sign a consent form. Please feel free to ask any questions that you have and remember that, even at this stage, you can decide against going ahead with the procedure if you so wish. You