



We can provide interpreters for a variety of languages, information in larger print or other formats (e.g. audio) - please call us on 01932 723553.

You can use Text Relay to turn telephone communications into text.  
Use the Relay UK app on your phone, tablet or PC.  
You can also use Relay UK via Minicom or Uniphone.

اگر نیاز به ترجمہ دارید، لطفاً با شماره 01932 723553 تماس بگیرید.

ਜੇ ਤੁਹਾਨੂੰ ਭਰਜਮੇ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਇਸ ਨੰਬਰ ਤੇ ਫੋਨ ਕਰੋ: 01932 723553

اگر آپ اس کا اردو زبان میں ترجمہ چاہتے ہیں، تو براہ کرم اس فون نمبر 01932 723553 پر رابطہ کریں

Se precisa de uma tradução por favor contacte: 01932 723553

আপনার অনুবাদের দরকার হলে এখানে যোগাযোগ করুন : 01932 723553

यदि आपको अनुवाद की ज़रूरत है तो कृपया इस नंबर पर फोन करें: 01932 723553

Jeżeli chcemy, aby te informacje w innym języku, proszę zadzwonić 01932 723553

**Ashford Hospital**  
London Road  
Ashford, Middlesex  
TW15 3AA  
Tel: **01784 884488**

**St. Peter's Hospital**  
Guildford Road  
Chertsey, Surrey  
KT16 0PZ.  
Tel: **01932 872000**

Website: [www.ashfordstpeters.nhs.uk](http://www.ashfordstpeters.nhs.uk)

## Percutaneous Drainage Interventional Imaging Department



# Percutaneous Drainage

## Patient Information

*Reproduced with the kind permission of the British Society of Interventional Radiology (BSIR) 2011*

This leaflet tells you about percutaneous drainage. It explains what is involved and what the possible risks are. It is not meant to replace an informed discussion between you and your doctor but can function as a starting point for such discussions. If you have any questions about the procedure, please ask the doctor who has referred you or the department which is going to perform it.

## What is percutaneous drainage?

In the past, drainage of an abscess inside your body would have required an open operation. Now it is possible to drain an abscess directly by inserting a thin plastic tube, called a drainage catheter, into it through the skin (percutaneous). This is typically performed through a small incision. This procedure is called percutaneous drainage.

## Why do you need drainage?

Imaging that you have had performed, such as an ultrasound scan or a CT scan, will have shown that you have an abscess that is suitable for draining through a small tube, rather than by having an

## Finally

Some of your questions should have been answered by this leaflet but remember that this is only a starting point for discussion about your treatment with the doctors looking after you. Make sure you are satisfied that you have received enough information about the procedure.

---

## Further Information

We endeavour to provide an excellent service at all times, but should you have any concerns please, in the first instance, raise these with the Matron, Senior Nurse or Manager on duty.

If they cannot resolve your concern, please contact our Patient Experience Team on 01932 723553 or email [asp-tr.patient.advice@nhs.net](mailto:asp-tr.patient.advice@nhs.net). If you remain concerned, the team can also advise upon how to make a formal complaint.

---

Author: Marites Lomibao

Department: Imaging

Version: 3

Published: Jan 2024

Review: Jan 2027

## **How long will it take?**

Every patient is different, and it is not always easy to predict; however, expect to be in the radiology department for about 30 minutes.

## **What happens afterwards?**

If you are inpatient, you will be taken back to your ward on a trolley. And, if you are a day case patient, you will be recovered in the imaging department until you are ready and safe to be discharged home. Nurses allocated to look after you will conduct routine observations, such as taking your pulse and blood pressure, to make sure that there are no problems. You will stay in bed for a few hours, until you have recovered. The drain will remain in the collection until it has been fully emptied. Once this has happened, the drain can be removed by the clinician or a trained nurse. This is not usually painful.

## **How long will the catheter stay in?**

These are questions which only the doctors looking after you can answer. It is possible that you will need further scans or X-rays to check that the collection has been drained completely. You will be able to lead a normal life with the catheter in place. When the catheter is taken out, this does not hurt at all.

open operation. Abscesses can make you ill, and if they occur after surgery, will delay your recovery. Although antibiotics can help, they cannot really be effective against a large abscess. However, once the pus has been drained, this can be sent to the laboratory for tests to show which the best antibiotic to treat the remaining infection is.

## **Are there any risks?**

Percutaneous drainage is a very safe procedure, but as with any medical procedure there are some risks and complications that can arise. The biggest problem is being unable to place the drainage tube satisfactorily into the abscess. If this happens, your consultants will arrange another method of draining the abscess, which may involve an operation. Rarely, you may get a shivering attack (a rigor) during the procedure, but this can be treated satisfactorily with antibiotics. Despite these complications, the procedure is normally very safe and will certainly result in a significant improvement in your medical condition.

## **Who has made the decision?**

The consultant in charge of your care and the interventional radiologist performing the procedure have discussed your case and feel that this is the best option. However, you will also have the opportunity for your opinion to be considered and if after discussion with your doctors, you no longer want the procedure, you can decide against it.

## **Are you required to make any special preparations?**

You will either be an inpatient in the hospital or a day case depending on the complexity of your condition. You will need to have a blood test performed beforehand to check that you do not have an increased risk of bleeding. You will have to be advised not to eat for four hours before the procedure, although you may still drink clear fluids such as water.

## **Who will you see?**

A specially trained team led by an interventional radiologist within the radiology department. Interventional radiologists have special expertise in reading the images and using imaging to guide catheters and wires to aid diagnosis and treatment.

## **Where will the procedure take place?**

In the radiology department - either in the ultrasound, CT scanner or a special X-ray room. It all depends on where the abscess is in the body and which imaging the radiologist feels is best for you.

## **What happens during the percutaneous drainage?**

This does depend on where the abscess is in your body and which imaging is being used. Usually, you will lie on your back or front in the position that the radiologist has decided is most suitable. The radiologist will explain this to you before performing the drainage.

You will be asked to get undressed and put on a hospital gown. You may receive a sedative to relieve anxiety. You may need to have a needle put into a vein in your arm, so that the radiologist can give you a sedative or painkillers. The procedure is performed under sterile conditions and the interventional radiologist and radiology nurse will wear sterile gowns and gloves to conduct the procedure. Your skin will be prepped with antiseptic, and you will be covered with sterile drapes. The radiologist will use an ultrasound probe, X-rays, or the CT scanner to decide on the most suitable point for inserting the drain. Local anaesthetic will be injected into the skin to numb the area. A fine needle is inserted into the abscess to obtain a sample. What happens next depends on what the sample looks like. Most of the time, a guide wire will be placed into the abscess to allow the correct positioning of a drainage tube (catheter). This will be connected to a drainage bag. Occasionally, the abscess may simply be drained through the needle or small plastic tube, which is withdrawn altogether.

## **Will it hurt?**

When the local anaesthetic is injected, it will sting for a short while, but this soon wears off. Some discomfort may be felt when the drain is placed.